

L# ANSWER 26 OF 419 CAPLUS COPYRIGHT 2002 ACS

AN 2001:288845 CAPLUS

DN 134:295735

TI Preparation of N-alkyl-2-pyrrolidone from .gamma.-butyrolactone or its
analogs

IN Takahashi, Kazushige; Seito, Yoko; Utsunomiya, Masaru; Orita, Soichi;
Oshu, Tomonori

PA Mitsubishi Chemical Corp., Japan

SO Jpn. Kokai Tokkyo Koho, 4 pp.

CODEN: JKXXAF

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

-----	----	-----	-----	-----
-------	------	-------	-------	-------

PI	JP 2001114760	A2	20010424	JP 1999-291972	19991014
----	---------------	----	----------	----------------	----------

AB N-alkyl-2-pyrrolidone is prepd. by treatment of .gamma.-butyrolactone (I),
4-hydroxybutyric acid (II), or its oligomer (III) with .gtoreq.1
alkylamine(s) in the presence of carboxylic acid (other than II or III),
carboxylate ester, or carboxamide. Thus, I was autoclaved with MeNH₂,
Me₂NH, Me₃N, and AcOMe in H₂O for 5 h to give 94.9%
N-methyl-2-pyrrolidone.

IC ICM C07D207-267

ICS B01J031-04; C07B061-00

DT Patent

LA Japanese